CFD Analysis of Loss Of Vacuum Accident for Safety Application

in Experimental Fusion Facility

C.Bellecci¹, P.Gaudio¹, <u>I.Lupelli¹</u>, A.Malizia¹, M.T.Porfiri², R.Quaranta¹, M.Richetta¹ ATOM-Faculty of Engineering, University of Rome "Tor Vergata", Via del Politecnico 1, 00133 Rome, Italy 2ENEA Nuclear Fusion Technologies, Via Enrico Fermi , 45 I-00044 Frascati, Italy



LOVA in ITER

The aim of the fusion research is to develop a prototype fusion power plant that is safe and reliable, environmentally responsible and economically viable. ITER will be the world's largest experimental facility that aims to demonstrate the scientific and technical feasibility of fusion power. In a fusion experimental reactor the dust is generated during normal machine operations and by macroscopic erosion of the plasma facing materials due to intense thermal loads that occur during Edge Localized Modes (ELMs), plasma disruptions (PD) and Vertical Displacement Events (VDEs). This dust, mainly accumulated closeness the divertor zone, can be mobilized outside the Vacuum Vessel (VV) in case of LOVA threatening public safety because it may contain tritium, may be radioactive and may be chemically reactive and/or toxic. A LOVA is a Design Basis Accident (DBA) event in fusion reactors [1]. Several studies on fundamental phenomena have been carried out in order to investigate the features of LOVA events both for the wet and dry case [2,3,4,5,6,7] and also reference events for ITER were postulated [8]. One of the main issue is to develop methods of computational analysis for the accident scenarios. An experimental facility, STARDUST, has been developed and utilized to validate new computational models capable of predicting the flow field for dust resuspension and transport in fusion reactor LOVA scenarios.





laist teaperature	Ti	204,31			Isothermal	Adiabatic	
Dated previous	34	180925			230122222		
Date: descrip	10	1,300	kg/m3	Pi(Pa)	112,9226	112.9226	
usity or conceined presents (200.3000E)	Cje	947,3	33060	PI(Pa)	112,9220	1112,9220	
with at countries volume (200-20008).	CHI	680	71085		-	,	
Salar Cp Cv venio	- 10	1,403		Pf(Pa)	101325	101325	
bler bysoak raconty distantia b	ų)	1,819-85	9n		101323	101323	
Sales Sciences in recording	10	1,024	mil h				
laint quad of small	val	346,0	m/s		294.15	294.15	
ble official previous	pot	59057		()	251,25	251,25	
Vend in Improve	Ti Ti	29(3)	K		294.15	418.254	
Yeard at radic previous	2)	112,9236			294,13	418,234	
Yeard air deploy	95	0,000,007	lg/m2				
mily of countries previous (200-1000E)	Cyc	947,3	31080		8.0650	8.0650	
(2000/-001) warder temano to etcor	01	800	31625	icuai(s)	0,0000	0,0000	
Venetial Cp.Cr ceta	96	1,403					
and six dynamic the suity (Sutherland)	16	1,618-45	Para	tc(s)	7.3133	5.1433	
Yeard Stansack recordy		1,018/42	823	10(3)	1,5155	5,2 155	
Vessel sir speed of smeat	100	3465	m/s	180	17 72 44	12 0004	
Venal sir critical pressure	pck	90	Ps.	tf(s)	11/./344	13.0084	

STARDUST Description

STARDUST is a stainless steel horizontal cylinder (internal volume 0.17 m3) closed by two lids [10] (Fig 1). STARDUST is equipped with an automatic data acquisition system that allows the control of internal pressure, wall temperature and air flow inlet in order to carry out the experiments at the desired initial conditions (see table 2). When the initial conditions are reached the control system opens the flow meter inlet valve and then the air flows inside the tank with a which the ingress occurs by pressure difference between the atmospheric condition and the internal condition of the tank (EX_B) without an inlet flow rate imposed. Air inlet simulates the loss of vacuum event . The air flow passes carried out introducing a semi cylindrical obstacle made of stainless steel, which simulates the presence of structures, in particular the divertor, inside the ITER VV [13,14]. In order to take into account the influence of the space between limiter and divertor, as in the VV of ITER, a slit has been realized on the obstacle (Fig 2). In the experimental campaign has been evaluated punctual flow velocity values, at 20cm from the inlet section, obtained in two different internal conditions: 1)Experiments without obstacle (VA_WO for A inlet and VB_WO for B inlet valve); 2)Experiments without obstacle in order to understand better the influence of structures (like divertor) present inside STARDUST facility (VA_O pressure Ps) and the pressure measured by the head of sensor (total pressure Pt). When the initial conditions inside the

has been used where γ is the ratio of the specific heat of the fluid at constant pressure to the specific heat of the fluid at constant volume, R is the universal gas constant, Tmean is mean temperature measured by internal thermocouples and M is the air molecular mass. The pressurization rate is measured at the top of the chamber with a Pirani sensor.

$$\begin{split} &\frac{\partial p}{\partial t} + \nabla \left(\rho u \right) = 0 \\ &\rho \frac{\partial u}{\partial t} + \rho u \cdot \nabla u = -\nabla p + \nabla \cdot \left[\eta \left(\nabla u + \left(\nabla u \right)^\intercal \right) - \left(\frac{2}{3} \eta - \kappa_{in} \right) \left(\nabla \cdot u \right) I \right] + F \end{split}$$

 $\rho C_p \left[\frac{\partial T}{\partial t} + (u \cdot \nabla)T \right] = -(\nabla \cdot q) + \tau : S - \frac{T}{\rho} \frac{\partial p}{\partial T} \Big|_p \left(\frac{\partial p}{\partial t} + (u \cdot \nabla)p \right) + Q$ $-n \cdot (-k\nabla T) = h(T_{\text{inf}} - T)$

Cp: is the specific heat capacity at constant pressure (J/(kg·K))
T: absolute temperature (K)
q: heat flux by conduction (W/m2)
t: viscous stress tensor (Pa)
S: strain rate tensor (1/s):
Q: contains heat sources other than viscous heating (W/m3)
h: heat transfer coefficient (W/(m2·K)).

$$\begin{split} t_{r,C}^{i} &= \left(\frac{2}{\gamma+1}\right)^{1/2} - P_{r,i} \left(\frac{2}{\gamma+1}\right)^{\frac{\gamma+1}{2(\gamma-1)}} & t_{r,C}^{i} &= \frac{1}{\gamma} \left(\frac{2}{\gamma+1}\right)^{1/2} - \frac{P_{r,i}}{\gamma} \left(\frac{2}{\gamma+1}\right)^{\frac{\gamma+1}{2(\gamma-1)}} \\ P_{r}^{i}(t_{r}) &= P_{r,i} + \left(\frac{2}{\gamma+1}\right)^{\frac{\gamma+1}{2(\gamma-1)}} t_{r} & P_{r}^{i}(t_{r}) &= P_{r,i} + \gamma \left(\frac{2}{\gamma+1}\right)^{\frac{\gamma+1}{2(\gamma-1)}} t_{r} \\ P_{r}^{j}(t_{r}) &= \left[1 - \left[\sqrt{1 - P_{r,c}^{\frac{j-1}{\gamma}}} - \sqrt{\frac{\gamma-1}{2\gamma^{2}}} (t_{r} - t_{r,c}^{j})\right]^{\frac{\gamma}{j-1}} \right] \end{split}$$

 $P_r^A(t_r) = \left[1 - \left[\sqrt{1 - P_{r,C}^{\frac{\gamma - 1}{\gamma}}} - \sqrt{\frac{\gamma - 1}{2}} \left(t_r - t_{r,C}^A\right)^2\right]^{\frac{\gamma}{\gamma}}\right]$

configurations characterizing the flow field is shown in figures 3 4 5 and 6.

in case of leakage at divertor ports level (valve B). This result matches with experiments done in 2007 experimental campaign [13,14] in which we have measured the quantity of dust resuspended in case of LOVA. The percentage of dust resuspended, that simulates a LOVA at divertor level is lower in case of obstacle inside the chamber [13,14]. These results values measured without obstacle (VB_WO) are higher than velocity values measured with obstacle (VB_O)

consol models for short transients (*1,5 s) have given reasonable results. Future work will focus on obtaining accurate results for longer transients, resolving flow features such as shocks and boundary layers, implementing of dust resuspension models, improving mesh and geometry impact on cpu time, including turbulence model behavior. The dust resuspension models include most of the transport phenomena relevant for ITER. The 3D numerical code describing the dust mobilization in STARDUST facility will be improved by means of COMSOL software and comparison will be made with experimental data acquired during 2007, in order to validate the model. Then the same model will be adapted to a toroidal shape and an optical diagnostic for measurement of dust mobilization evolution will be chosen and studied. The results will be useful for the design of optical ports in a new experimental facility with a geometrical shape similar to ITER foreseen to be realized.









